


Pr addyi[®]
(flibanserin)
100mg tablets

Getting started with Pr ADDYI[®]



ADDYI (flibanserin) is used to treat a condition called hypoactive sexual desire disorder (HSDD) in women. This means that you have had low sexual desire for a minimum of 6 months that happens 75–100% of the time and this causes you distress or difficulty in your relationships. The low sexual desire must not be a result of some other condition (medical or psychological), problems in a relationship, or effects of a medication or other drug substance.

It is used in pre and naturally postmenopausal women, aged 60 years or younger, who:

- have not had problems with low sexual desire in the past, and
- have low sexual desire regardless of the type of sexual activity, situation, or sexual partner.

What is hypoactive sexual desire disorder (HSDD)?

HSDD is a condition marked by low sexual desire for a minimum of 6 months that happens 75–100% of the time and may cause you distress or difficulty in your relationship.

HSDD has been medically recognized for nearly half a century and has affected many just like you – you are not alone



HSDD is the **most common** form of female sexual dysfunction.



In HSDD, the low sexual desire persists no matter the type of sexual activity, the situation, or the sexual partner.

Why ADDYI?

- ADDYI is used for the treatment of pre and naturally postmenopausal women, aged 60 years or younger, with HSDD.
- It is prescribed when a woman has had low sexual desire, 75–100% of the time, for a minimum of 6 months, and when the situation causes her difficulty in her relationship.
- ADDYI does not improve sexual performance.
- Before prescribing, your healthcare professional may have used a checklist that includes a number of specific questions to make sure that ADDYI is right for you.

This is Addyi®  A little Pink Pill*

* Not actual size

How to take ADDYI

- Take ADDYI exactly as your healthcare professional tells you to: **one tablet, once-a-day, and only at bedtime.**
- Do not take more than the dose prescribed by your healthcare professional.
- Never take ADDYI at another time of the day, otherwise you may
 - get low blood pressure,
 - faint (lose consciousness), and/or
 - get very sleepy.
- If you miss a dose of ADDYI, simply take your next dose at bedtime the next day. Do not take ADDYI the next morning. Do not double your next dose.



Talk to your healthcare professional if your symptoms of HSDD have not gotten better after taking ADDYI for 8 weeks

Serious warnings and precautions

- ADDYI can cause severe low blood pressure and fainting.
- Take ADDYI at bedtime. If you take ADDYI and feel lightheaded or dizzy, lie down right away. Drinking alcohol close to the time you take ADDYI increases your risk of severe low blood pressure and fainting.
- ADDYI can cause sedation or make you feel very drowsy. After taking ADDYI, do not drive or perform other activities that require full attention until you know how it affects you. Using ADDYI with depressant drugs may increase the risk of sedation or drowsiness.
- If you have liver problems, you should not take ADDYI.
- ADDYI should not be taken if you are taking certain other medicines.
- Before taking ADDYI, tell your healthcare professional about all the medicines you take, including drugs, vitamins, minerals, natural supplements, or alternative medicines. They will tell you if it is safe to take ADDYI.

Other warnings you should know about

- The effects of ADDYI were not studied in Black or Asian women who are postmenopausal, or postmenopausal women who are underweight.
- If you weigh less than 50 kg, there might be a higher risk of certain side effects, including central nervous system depression (sedation or feeling very sleepy), low blood pressure and fainting. Tell your healthcare professional right away if you feel these effects.
- If you drink alcohol, wait at least 2 hours after drinking 1 or 2 standard alcoholic drinks* before taking ADDYI at bedtime. If you drink 3 or more standard alcoholic drinks in the evening, skip your ADDYI dose at bedtime. After you have taken your ADDYI at bedtime do not drink alcohol until the following day.
- ADDYI can cause low blood pressure, fainting (loss of consciousness) and sleepiness. This can affect your ability to drive or use machines. You should wait at least 6 hours after taking ADDYI before driving or using machines. You should be extra careful when driving the morning after taking ADDYI until you know how it affects you.
- Do NOT drink grapefruit juice when taking ADDYI.

* Examples of 1 standard alcoholic drink include: one 12-ounce regular beer (5% alcohol); 5 ounces of wine (12% alcohol); 1.5 ounces of distilled spirits or shot (40% alcohol).




Possible side effects

Just like with any medication, you may experience certain side effects while taking ADDYI. Some side effects may include:

- dizziness
- vertigo
- nausea, vomiting
- muscle spasms
- swelling of hands and legs
- sleepiness, tiredness
- night sweats
- difficulty falling asleep or staying asleep
- dry mouth
- anxiety
- diarrhea
- constipation
- indigestion
- abdominal pain
- headache
- abnormal dreams
- rash
- acne

**This is not a complete list of side effects.
For any unexpected side effects, talk to your
healthcare professional.**

Please refer to the Patient Medication Information section of the ADDYI Product Monograph for more information.



You Are
Not Alone



Pr **addyi**[®]
(flibanserin)
100mg tablets

Let's talk about
the elephant
in the bedroom!



For more information on ADDYI and
HSDD, use the DIN located on your
ADDYI bottle to access addyi.ca or
talk to your healthcare professional.

SEARCHLIGHT
PHARMA

Searchlight Pharma Inc.
1600 Notre Dame St. West, Suite 312
Montréal, Québec H3J 1M1
www.searchlightpharma.com

ADDYI is a registered trademark.
© Searchlight Pharma Inc. 2021
Printed in Canada – April 2021

